
Effect of *Mucuna pruriens* leaves in dairy cow feed on gas production, digestibility and rumen fermentation by using *in vitro* gas production technique

Yeanpet, C. *, Pilajun, R., Lunpha, A. and Glinubon, J.

Department of Animal Science, Faculty of Agricultural, Ubon Ratchathani University, Ubon Ratchathani, Thailand.

Yeanpet, C., Pilajun, R., Lunpha, A. and Glinubon, J. (2022). Effect of *Mucuna pruriens* leaves in dairy cow feed on gas production, digestibility and rumen fermentation by using *in vitro* gas production technique. International Journal of Agricultural Technology 18(3):1331-1338.

Abstract The results showed that the *Mucuna pruriens* leaves at 10% in feed affect dry matter and organic matter digestibility which higher than with the control group ($p < 0.05$). However, *M. pruriens* leaves in all level were not different among dietary treatments on the gas production from the immediately soluble fraction, the gas production from the insoluble fraction, and the gas production rate constant for the insoluble fraction and the potential extent of gas production. Therefore, it can use *M. pruriens* leaves at 10% in dairy cow feed.

Keywords: *Mucuna pruriens*, Gas production, Digestibility, Rumen fermentation

Introduction

Dairy farming is an important occupation in Thailand because people tend to increase milk consumption and milk is widely as a nutritious drink for people of all ages. Especially, among children (Zhang *et al.*, 2021). In addition, the Thai government has the policy to supported farmers to enhance dairy production. However, farmer has still faced many issues in the dairy production sector such as management, disease, breed and feed. Especially, feed costs have an increased. Most of the farmers in Thailand depend on natural pastures but the roughages are poor quality and insufficient of roughage for animal so require protein supplementation (Muinga and Saha, 2003) but the ingredient as protein source in concentrate was *expensive* such as soybean meal, Palm kernel meal, fish meal and cottonseed meal (Abdeltawab and Khattab, 2018). Agricultural residues include rice straw, wheat straw and corn stover, which are used as animal feed. In addition, native plant was used to feed animals, it consists of *Leucaena*, *Mata Raton*, *Mulberry* and *Mucuna*.

* Corresponding Author: Yeanpet, C.; Email: chittraporn.y@ubu.ac.th

Mucuna (*Mucuna pruriens*) is widespread in tropical and sub-tropical regions of the world and with a great potential for animal feeding. This is the high protein content which can be used as a replacement for soybean meal in animal feeding. The chemical and nutritional composition of *Mucuna* contains 25-35% crude protein, 31.2-39.5 % of starch and 21.3 % neutral detergent fiber (Martínez-Pérez *et al.*, 2008). Several research studies have been carried out on the development about using *M. pruriens* leaves in animal feed to increase the use of alternative protein sources and reduce feed cost. According to Muir and Saha (2003) who found that cows fed *Mucuna* forage tended to have increased production performance and live bodyweight and *Mucuna* hay has been fed both to dairy cattle to increase milk production and meat production (Muthia and Gachui, 2003). However, the studies and research about *Mucuna pruriens* leaves in dairy cow feed is limited and unclear. Therefore, the objective of this study was to investigate the effect of *M. pruriens* leaves meal as a protein source in dairy cow feed on gas production, digestibility and rumen fermentation by using *in vitro* gas production technique.

Materials and methods

Study area

The study was carried out in Ubon Ratchathani University, Ubon Ratchathani Province which is located in the north-eastern region of Thailand during June 2020- June 2021.

Experimental design and dietary treatments

The experiment was performed in a Completely Randomized Design (CRD). The basal diet was a 40:60 ratio of Ruzi grass (*Brachiaria ruziziensis*) and concentrate. The ingredients composition of the basal diet is shown in Table 1. There were four dietary treatments consisting of basal diets without *Mucuna pruriens* (the control group; T1) and T2, T3, T4 were basal diets using *M. pruriens* at 10, 20 and 30% respectively. Dietary treatments were incubated at 70 °C for three days and ground into a powder form and passed through a 0.5 mm sieve.

Rumen and substrate inoculum

Rumen fluid was collected from three healthy dairy cows, which were placed on a routine basis for at least 1 week and fed by the same basal diet for 5

days before sampling by collection through a suction pump. The rumen fluid was collected and filtered through two layers of cheesecloth. *In vitro* fermentation was used in this study according to the technique described by Makkar *et al.* (1995). The rumen fluid (660 ml) was added to warm (39 °C) and reduced medium consisting of 1,095 ml distilled water, 730 ml rumen buffer solution (417 mM NaHCO₃ and 51 mM NH₄HCO₃), 365 ml macro mineral solution (46 mM KH₂PO₄, 40 mM Na₂HPO₄, 38 mM NaCl and 2 mM MgSO₄•7H₂O), 0.23 ml micromineral solution (505 mM MnCl₂•4H₂O, 898 mM CaCl₂•2H₂O, 42 mM CoCl₂ • 6H₂O and 341 mM FeCl₂ • 6H₂O), 1 ml of 4mM resazurin and 60 ml freshly prepared reduction solution (145 mM Na₂S•9H₂O and 3.7 ml 1 M-NaOH). The mixture was kept stirring and continuously filled with CO₂ to ensure anaerobic condition at 39 °C on a hot plate. Approximately 30 ml of the rumen-fluid medium was transferred into serum bottles incubated with dietary treatments (200 mg) at 39 °C for 72 h.

Table 1. Ingredient composition of the basal diet

Item	% of fresh weigh
Ingredients	
Maize	25.0
Cassava chip	22.0
Rice bran	10.0
Soybean meal	26.4
Palm kernel meal	15.0
Mineral premix	0.5
Salt	0.5
sulfur	0.1
Dicalcium	0.5

Data collection and analysis

Dietary treatments were analyzed for dry matter (DM), Ash, crude protein (CP) and ether extract (EE) using the procedure of AOAC (1997), while neutral detergent fiber (NDF) and acid detergent fiber (ADF) were also determined as described by Van Soest *et al.* (1991).

Gas production during the incubation, the kinetics of gas production were recorded at 2, 4, 6, 8, 12, 18, 24, 36, 48 and 72 h. Cumulative gas production data were fitted to the model of Ørskov and McDonald (1979) as follows:

$$Y = a + b (1 - e^{-ct})$$

Where a = the gas production from the immediately soluble fraction, b = the gas production from the insoluble fraction, c = the gas production rate constant for the insoluble fraction (b), t = incubation time, (a+b) = the potential extent of gas production, y = gas production at time 't'. At 24- and 48-h post inoculation, a set of samples were determined *in vitro* true digestibility according to Van Soest and Robertson (1985).

Statistical analysis

All data were analyzed as a Completely Randomize Design (CRD) using the SAS (1998). The significant differences between treatments were compared by Duncan's New Multiple Range Test (Steel and Torrie, 1980). Difference among treatment with P<0.05 was accepted as statistical differences.

Results

The chemical compositions of basal diet and *Mucuna pruriens* leaves is shown in Table 2. The results showed the crude protein content in basal diet and *M. pruriens* leaves were 19.2% and 24.3% and 4.5% and 2.4% EE of dry matter basis.

Table 2. The chemical composition of basal diet and *Mucuna pruriens* leaves

Item	Basal diet	<i>Mucuna pruriens</i> leaves
DM (%)	96.5	17.4
Chemical composition		
OM	91.8	94.6
Ash	8.2	5.40
CP	19.1	24.3
EE	4.5	2.40
NDF	32.2	66.2
ADF	27.5	44.8

DM = Dry Matter, OM = Organic Matter, CP = Crude Protein, EE = Ether Extract
NDF = Neutral Detergent Fiber, ADF = Acid Detergent Fiber

Kinetic of gas production

The estimated parameters of gas production are shown in Table 2 and cumulative gas production profiles are shown in Figure 1. The results showed

that the gas production from the insoluble fraction (b) the potential extent of gas production (d) and effective gas production potential (EP) were significant different among treatments ($P < 0.05$), of which the *Mucuna pruriens* leaves at 30% that was lower than with the control group and *M. pruriens* leaves at 10%. However, the gas production from the immediately soluble fraction (a), the gas production from the insoluble fraction (b), the gas production rate constant for the insoluble fraction (c) were not significant different among treatments ($P > 0.05$). However, the gas production at different incubation times (GP 2, 4, 6, 8, 12, 18, 24, 36, 48 and 72 h) (Figure 1).




Figure 1. Effect of *Mucuna pruriens* leaves on gas production at different times of incubation: T1 = Basal diet (Control group), T2 = Basal diet used *Mucuna pruriens* leaves at 10%, T3 = Basal diet used *Mucuna pruriens* leaves at 20%, T4 = Basal diet used *Mucuna pruriens* leaves at 30%

Table 3. Effects of *Mucuna pruriens* leaves on gas production in dietary treatments

Item	Dietary treatments				SEM	P-value
	T1	T2	T3	T4		
Gas production characteristics						
a	-1.718	-1.755	-1.320	-1.775	0.13	0.61
b	55.2 ^a	55.4 ^a	51.9 ^{ab}	48.8 ^b	1.01	0.04
c	0.0682	0.0681	0.0686	0.0699	0.001	0.88
d	53.5 ^a	53.6 ^a	50.5 ^b	47.0 ^c	0.96	0.03
Effective gas production potential (EP)						
	30.1 ^a	30.1 ^a	28.7 ^{ab}	26.6 ^b	0.49	0.02

The significant at P < 0.05

a = the gas production from the immediately soluble fraction, b = the gas production from the insoluble fraction, c = the gas production rate constant for the insoluble fraction, d = the potential extent of gas production, EP = $a + [bc/(k + c)]$ where $k = 0.05$ (Ørskov and McDonald, 1979)

Table 4. Effects of *Mucuna pruriens* leaves on *in vitro* digestibility

Items	Dietary treatments				SEM	P-value
	T1	T2	T3	T4		
<i>In vitro</i> dry matter digestibility, IVDMD (%)						
24 h	45.5 ^a	46.0 ^a	40.4 ^b	40.0 ^b	1.07	<0.01
48 h	52.5 ^a	56.1 ^a	46.9 ^b	47.0 ^b	1.51	<0.01
<i>In vitro</i> organic matter digestibility, IVOMD						
24 h	47.7 ^b	55.3 ^a	47.3 ^b	41.5 ^c	1.87	<0.01
48 h	63.5 ^b	65.5 ^a	59.4 ^c	58.8 ^d	1.05	<0.01
Metabolizable energy, ME (Mcal/kgDM)						
24 h	2.07 ^a	2.08 ^a	2.01 ^{ab}	1.92 ^b	0.02	0.03
48 h	2.38 ^a	2.39 ^a	2.29 ^{ab}	2.18 ^b	0.03	0.02
Total digestible energy, TDN						
24 h	56.7 ^a	56.7 ^a	55.3 ^{ab}	53.3 ^b	0.52	0.04
48 h	63.3 ^a	63.7 ^a	61.5 ^{ab}	59.1 ^b	0.66	0.02

Significant at $P < 0.05$, Significantly at $P < 0.01$

T1 = Basal diet (Control group), T2 = Basal diet used *Mucuna pruriens* leaves at 10%,

T3 = Basal diet used *Mucuna pruriens* leaves at 20%, T4 = Basal diet used *Mucuna pruriens* leaves at 30%

In vitro organic matter digestibility (IVOMD) in basal diet used *M. pruriens* leaves at 10% (T2) was higher than the other groups ($P < 0.01$). Moreover, *in vitro* dry matter digestibility was not affected between the control group and basal diet used *M. pruriens* leaves at 10% but basal diet used *M. pruriens* leaves at 20% and 30% were lower than with the control group ($P < 0.01$). Metabolizable energy (ME) and total digestible energy (TDN) incubated at 24 h and 48 h were significant between dietary treatments ($P < 0.05$) by the basal diet used *Mucuna pruriens* leaves at 30% was lower the other group. However, basal diet used *M. pruriens* leaves at 10, 20% was no different with control group.

Discussion

The crude protein content in *M. pruriens* leaves is and 24.3% and 4.5% EE of dry matter basis. This results similarly reported by Ujowundu (2010) who reported the *M. pruriens* leaves contain 25% crude protein of dry matter basis. Basal diet was also formulated to contain 22.1% CP, 4.5% EE, 97.8% OM, 32.2 % NDF and 27.5% ADF of dry. The results of the gas production

from the immediately soluble fraction (a) were -1.718, -1.755, -1.320 and -1.775 respectively that result was minus because the treatment diets are resting state. This result agrees with Yeanpet *et al.* (2021). The basal diet used *M. pruriens* leaves at 30% affected on the gas production from the insoluble fraction (b) the potential extent of gas production (d) and effective gas production potential (EP) lower than with the control group and basal diet used *M. pruriens* leaves at 10% that *M. pruriens* leaves contain 12.5% crude fiber of dry matter basis (Ezeokonkwo and Okafor, 2015), whereas soybean meal used as a source of protein in basal diet has higher fiber content ($\approx 5\%$ of DM) (Stein *et al.*, 2008). The high fiber content in the dietary treatments effect on digestion and kinetic of gas production. However, there are only a few studies about *M. pruriens* leaves on digestibility in cow. While *M. pruriens* leaves at 10% no different with control group while the gas production from the immediately soluble fraction (a), the gas production from the insoluble fraction (b), the gas production rate constant for the insoluble fraction (c) were not significant difference among treatment.

The *in vitro* fermentation profiles were significantly different of *Mucuna pruriens* leaves for *in vitro* organic matter digestibility (IVOMD) at 24 and 48 h. *Mucuna pruriens* leaves at 10% higher than with another group but *M. pruriens* leaves at 30% lower than with other group this result suggests that *M. pruriens* leaves consist polyphenols, trypsin inhibitors, phytate, saponins, lectins, and tannin able to bind to proteins, thus lowering their digestibility. (Lampariello *et al.*, 2011). *Mucuna pruriens* at 30% had negative effect on metabolizable energy (ME) and total digestible energy (TDN) at 24 h and 48 h because *in vitro* dry digestibility in *M. pruriens* leaves at 30% lower than another group. While basal diet used *Mucuna pruriens* at 10, 20% was no different with control group (Muinga and Saha, 2003)

This study shown the results that the *M. pruriens* leaves at 10% enhance *in vitro* dry matter and organic matter digestibility and which has not a negative impact of kinetic of gas production of cow can use *M. pruriens* leaves at 10% in dairy cow feed. However, the issue regarding *M. pruriens* leaves supplement in feed on digestibility has not yet been completely settled. Further studies are warranted to understand these relationships

Acknowledgements

The authors would like to express their most sincere gratitude to The Ubon Ratchathani University for financially support this research project and Faculty of Agriculture, Ubon Ratchathani University for the use of research facilities.

References

- Abdeltawab, A. M. and Khattab, M. S. A. (2018). Utilization of palm kernel cake as a ruminant feed for animal: A review. *Asian journal biological sciences*, 11:157-164.
- AOAC. (1997). *Official Method of Analysis*. 16th Ed., Association of Official Agriculture Chemicals, Washington, DC. USA.
- Ezeokonkwo, M A. and Okafor, S. N. (2015). Proximate Composition and Mineral Analysis of *Mucuna utilis* (Velvet Bean). *IOSR Journal of Applied Chemistry*, 8:42-45.
- Lampariello, L. R., Cortelazzo, A., Guerranti, R., Sticozzi, C. and Valacchi, G. (2011). The Magic Velvet Bean of *Mucuna pruriens*,” *Journal of Traditional and Complementary Medicine*, 1:331-339.
- Makkar, H. P. S., Blummel, M. and Becker, K. (1995). Formation of complexes between polyvinyl pyrrolidones or polyethylene glycols and tannins, and their implication in gas production and true digestibility in in vitro techniques. *British Journal of Nutrition*, 73:897-913.
- Martínez-Pérez, M., Sarmiento-Franco, L. A. Santos-Ricalde, R. H. and Sandoval-Castro, C. A. (2018). Chemical composition and in vitro protein digestibility of processed *Mucuna pruriens*. *Cuban Journal of Agricultural Science*, 52:313-319.
- Mbuthia, E. W. and Gachui, C. K. (2003). Effect of inclusion of *Mucuna pruriens* and *Dolichos lablab* forage in Napier grass silage quality and on voluntary intake and digestibility in sheep. *Tropical and Subtropical Agroecosystems*, 1:123-128.
- Muinga, R. W. and Saha, H. M. (2003). The effect of *Mucuna* (*Mucuna pruriens*) forage on the performance of lactating cows. *Tropical and Subtropical Agroecosystems*, 1:87-91.
- Ørskov, E. R. and McDonald, I. (1979). The Estimation of Protein Degradability in the Rumen from Incubation Measurements Weighted According to Rate of Passage. *The Journal of Agricultural Science*, 92:499-503.
- SAS (1998). *User's Guide: Statistic*, Version 6, 12th ed. SAS Inst. Inc., Cary, NC.
- Steel, R. G. D. and Torrie, J. H. (1980). *Principles and Procedures of Statistics*. McGraw Hill Book Co., New York: USA.
- Ujowundu, C. o., Kalu, F. N., Emejulu, A. A., Okafor O. E., Nkwonta, C. G. and Nwosunjoku E. C. (2010) Evaluation of the chemical composition of *Mucuna utilis* leaves used in herbal medicine in Southeastern Nigeria. *African Journal of Pharmacy and Pharmacology*, 4:811-816.
- Van Soest, P. J. and Robertson, J. B. (1985). *Analysis of Forages and Fibrous Foods*. A Laboratory Manual for Animal Science. Department of Animal Science, Cornell University, Ithaca, NY: USA.
- Van Soest, P. J., Robertson, J. B. and Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74:3583-3597.
- Yeanpet, C., Thamrongyoswittayakull, C., Wachirapakorn, C. and Songsermsakul, P. (2021). The efficacy of yeast cell wall in the adverse effect of zearalenone on ruminal fibrolytic bacteria and in vitro fermentation. *Songklanakarin Journal of Science and Technology*, 43:623-629.
- Zhang, X., Chen, X., Xu, Y., Yang, J., Du, L., Li, K. and Zhou, Y. (2021). Milk consumption and multiple health outcomes: umbrella review of systematic and meta-analysis in humans. *Nutrition & Metabolism*, 18:1-18.

(Received: 15 August 2021, accepted: 10 April 2022)